

71°



**SOCIETÀ ITALIANA DELLE
SCIENZE VETERINARIE**

CONVEGNO SISVET

In collaborazione con

Dipartimento
Medicina Veterinaria
Produzioni Animali



Istituto Zooprofilattico
Sperimentale
del Mezzogiorno



"La baia di Napoli" Renoir, 1881

XVII Convegno SICV

XV Convegno SIRA

XIV Convegno AIPVET

XII Convegno SOFIVET

IV Convegno RNIV

I Convegno ANIV

28 Giugno - 1 Luglio 2017

Università degli Studi di Napoli "Federico II"

Corso Umberto I, 40 - 80138 - Napoli

I contributi presenti negli Atti del 71° Convegno SISVet 2017

potranno essere citati utilizzando il codice ISBN 9788890909245

DETECTION OF *Streptococcus equi* DNA IN *Rhipicephalus bursa* TICKS FROM SOUTHERN ITALY

Capone A.¹, Cappelli A.¹, Valzano M.¹, Bozic J.¹, Preziuso S.¹, Marcoccia R.¹, Mensah P.¹, Chiodera A.², Epis S.³, Rinaldi L.⁴, Cringoli G.⁴, Favia G.¹, Ricci I.¹

¹School of Biosciences and Veterinary Medicine, University of Camerino, Camerino, Italy. ²Infectious Diseases Unit, Hospital of Macerata, Macerata, Italy. ³Department of Veterinary Science and Public Health, University of Milan, Milan, Italy. ⁴Department of Veterinary Medicine and Animal Production, University of Naples, Naples, Italy.

Ticks are obligated ecto-parasites of wild and domestic animals and occasionally humans. They have veterinary interest because vectors of different viruses, bacteria and protozoa. Among the most common species in Italy and Europe there are *Ixodes ricinus* and *Rhipicephalus* spp. The tick-borne diseases epidemiologically relevant in our country are: Lyme disease, rickettsiosis, relapsing fever, tularemia, meningoencephalitis and ehrlichiosis.

Recently, the extensive characterization of the tick-associated microbiome faces the possible role of ticks in the transmission of additional pathogens, which are known to be transmitted by other arthropod vectors. Through the Denaturing Gradient Gel Electrophoresis (DGGE), we analysed the bacterial community of seven ticks belonging to three species from different areas of Italy: no. 2 *Rhipicephalus sanguineus* (Lombardia), no. 2 *Rhipicephalus bursa* (Campania) and no. 3 *Ixodes ricinus* (Marche). The samples were analysed selecting the highly variable V3 region of the bacterial 16S rRNA gene as target. The dominant DGGE bands were purified and sequenced allowing the identification of the bacteria present in each samples.

Burkholderia sp., *Coxiella*-like endosymbiont, unc. *Clostridium*, *Rickettsia peacockii*, unc. *Staphylococcus* and *Xanthomonas* sp. were detected, as already described in ticks. Moreover, *Streptococcus equi* subsp. *zooepidemicus/ruminatorum* was identified in *R. bursa* collected from buffalos living in rural areas of Southern Italy (the high genetic identity does not allow the distinction of these subspecies, that is usually achieved by biochemical tests on cultured bacteria). The DGGE outcome was confirmed by multiplex-PCR in other *R. bursa* samples collected from buffalos, ponies and goats from the same area. In particular, no. 6 positive samples over no. 15 analyzed were detected (no. 2 from buffalos and no. 4 from ponies).

S. equi is a complex-species including *S. e. equi*, *S. e. zooepidemicus* and *S. e. ruminatorum*: *S. e. equi* is the ethological agent of strangles, a highly contagious and serious infection of horses, whereas *S. e. zooepidemicus* and *S. e. ruminatorum* can cause several infections in both animals and humans. All these pathogens are known to be transmitted through direct contact.

The DNA detection of *S. e. zooepidemicus/ruminatorum* in *R. bursa* is particularly interesting since to our knowledge this pathogen was not previously detected in any tick species. Thus, in-depth studies are worthy to assess an effective competence of ticks as vectors of *S. equi* subspecies, since if a potential role of *R. bursa* in their transmission will be confirmed, new perspectives in the control of these zoonoses will be opened.

[1] Dantas-Torres et al. (2011) Parasit Vectors 4:56. [2] Rudolf et al. (2009) Folia Microbiol 54:419-28. [3] Pisoni et al. (2009) J Dairy Sci. 92:943-51. [4] Preziuso and Cuteri (2012) J Equine Vet Sci. 32:292-96. [5] Marchandin et al. (2007) Emerg Infect Dis. 13:1964-66. [6] Fulde and Valentin-Weigand (2013) P. Curr Top Microbiol Immunol. 368:49-81.

Acknowledgments: The research leading to these results has received funding from the EU7°FP under the GA n°281222 to IRENE RICCI for the "SymbioVec" project.