

Supporting Information to the paper Biurrun I. et al. Benchmarking plant diversity of Palaearctic grasslands and other open habitats. *Journal of Vegetation Science*.

Appendix S7. Funding information.

GrassPlot development has been supported by the Bavarian Research Alliance (BayIntAn_UBT_2017_58), the Eurasian Dry Grassland Group (EDGG) and the International Association for Vegetation Science (IAVS); IB, CorM, JAC, IGM, DGM, MHe, DL and MTo were supported by the Basque Government (IT936-16); CorM, IAx, MCh, JDa, PD, MHá, ZL, ZPr, EŠ and LT were supported by the Czech Science Foundation (19-28491X); TR was supported by the Estonian Research Council (PUT1173); RJP was funded by the Strategic Research Programme of the Scottish Government's Rural and Environmental Science and Analytical Services Division"; SBa was supported by the GINOP-2.3.2-15-2016-00019 project; GFi was partially supported by the MIUR initiative "Department of excellence" (Law 232/2016); BJA was funded by the Spanish Research Agency (grant AEI/10.13039/501100011033); AK, VB, IM, DS, IV and DV were supported by the National Research Foundation of Ukraine (2020.01/0140); MP and AH were supported by the Estonian Research Council (PRG874, PRG609), and the European Regional Development Fund (Centre of Excellence EcolChange); Data collection of HCP was funded by FORMAS (Swedish Research Council for Environment, Agricultural Science and Spatial Planning) and The Swedish Institute; JR was supported by the Czech Science Foundation (grant No. 20-09895S) and the long-term developmental project of the Czech Academy of Sciences (RVO 67985939); ATRA was funded by the Grant of Excellence Departments, MIUR-Italy (ARTICOLO 1, COMMI 314 – 337 LEGGE 232/2016); JMA was supported by Carl Tryggers stiftelse för vetenskaplig forskning and Qatar Petroleum; AAli was supported by the Jiangsu Science and Technology Special Project (Grant No. BX2019084), and Metasequoia Faculty Research Startup Funding at Nanjing Forestry University (Grant No. 163010230), and he is currently supported by Hebei University through Faculty Research Startup Funding Program; ZB was supported by the NKFI K 124796 grant; The GLORIA-Aragón project of JLBA was funded by the Dirección General de Cambio Climático del Gobierno de Aragón (Spain); MCs and LDem were supported by DG Environment through the European Forum on Nature Conservation and Pastoralism and Barbara Knowles Fund, in collaboration with Pogány-havas Association, Romania; JDa was partially supported by long-term research development project no. RVO 67985939 of the Czech Academy of Sciences; BD and OV were supported by the NKFI KH 126476, NKFI KH 130338, NKFI FK 124404 and NKFI FK 135329 grants; BD, OV and AKe were supported by the Bolyai János Scholarship of the Hungarian Academy of Sciences; BE was funded by the Environmental Department of the Tyrolean Federal State Government, the MAB Programme of the Austrian Academy of Science, the Mountain Agriculture Research Unit and the Alpine Research Centre Obergurgl of Innsbruck University. The GLORIA projects of BE were funded by the EU project no. EVK2-CT-2000-00056, the Earth System Sciences Program of the Austrian Academy of Sciences (project MEDIALPS), the Amt für Naturparke, Autonome Provinz Bozen-Südtirol, the Südtiroler Wissenschaftsfonds and the Tiroler Wissenschaftsfonds; RGG was supported by the Spanish Ministry of Research to sample GLORIA sites in central Spain (CGL 2008-00901/BOS) and present works by the Autonomous Region of Madrid (REMEDINAL TE-CM, S2018/EMT-4338); MJ was supported by National Geographic Society (NGS-288R-18) and Slovak Academy of Sciences (VEGA 2/0095/19); Alo was supported by national funds through FCT – Fundação para a Ciência e a Tecnologia (DL57/2016/CP1440/CT0001); ANo was partially funded by the National Science Centre, Poland, grant no. 2017/25/B/NZ8/00572; VO thanks Russian Science Foundation (#19-14-00038); SR was supported by the University of Latvia grant Nr. AAp2016/B041//Zd2016/AZ03; VS was supported by FCT-Foundation for Science and Technology PhD Programme SUSFOR (grant PD/BD/52696/2014);

CT was supported by the NKFI PD 132131 grant; YW and KW were supported by German Federal Ministry of Education and Research within the joint research program "Central Asia – Monsoon dynamics and geoecosystems (CAME) and by the German Ministry for Science, Research and Education under the BMBF - CAME Programm, project PaDeMoS (No. 03G0808A), work of KW and colleagues in Mongolia was supported by the German Science Foundation - DFG (WE 2601/3-1, 3-2) and by FWF - Der Wissenschaftsfonds, work in Yakutia again through German Science Foundation - DFG (WE 2601/9-1); Data collection by the Norwegian Institute of Bioeconomy Research was funded by the Ministry of Agriculture and Food through the 3Q Programme, whilst analyses and writing were funded by Research Council of Norway Grant No. 194051; NP and SŠ were partly supported by the Slovenian Research Agency, core fundings P1-0403 and J7-1822.